

Inflationary Cosmology in RS-I

Michele Ferraz Figueiró*

*Instituto de Física, Universidade de São Paulo
C.P. 66318, 05315-970, São Paulo-SP, Brazil*

Abstract

In this work, I intend to show a possible candidate of inflaton potential $V(\phi)$ in a scenario of a brane world defined by a pair of branes (RS-I).

PACS numbers:

The *Inflationary Cosmology* describes a phase in which our Universe evolves through accelerated expansion in a short time period at high energy scales. During this phase, our Universe is dominated by a potential $V(\phi)$ generated by a homogeneous scalar field $\phi(t)$ called *inflaton*. This potential must obey *slow-roll conditions* $\{\epsilon, |\eta| \ll 1\}$ where ϵ and η are the *slow-roll parameters*. These parameters are given by [1]

$$\epsilon(\phi) = \frac{M_{PL}^2}{2} \left(\frac{V'}{V} \right)^2 \quad (1)$$

and

$$\eta(\phi) = M_{PL}^2 \frac{V''}{V}. \quad (2)$$

We can calculate the spectral index $n(\phi)$ and its derivate for this potential $V(\phi)$

$$n - 1 = -6\epsilon + 2\eta \quad (3)$$

and

$$\frac{dn}{d \ln k} = -16\epsilon\eta + 24\epsilon^2 + 2\zeta \quad (4)$$

where

$$\zeta = M_{PL}^4 \frac{V'V'''}{V^2}. \quad (5)$$

The amount of inflation that occurs is described by the *number of e-foldings* N , is given by

$$N \equiv \ln \frac{a(t_{end})}{a(t)} \equiv \int_t^{t_{end}} H dt \approx \frac{1}{M_{PL}^2} \int_{\phi_{end}}^{\phi} \frac{V}{V'} d\phi, \quad (6)$$

where ϕ_{end} is defined by $\epsilon(\phi_{end}) = 1$ if inflation ends through violation of the slow-roll conditions.

The *Brane Cosmology* describes cosmological models with extra dimensions. A lot of interest in brane cosmology arose with a publication of two papers by *Randall* and *Sundrum* in the 90s. They propose a new higher-

dimensional mechanism for solving the hierarchy problem building two models, *RS-I* [2] and *RS-II* [3]. In these two models, they consider that the Standard Model particles and forces, with exception of gravity are confined to a *four-dimensional subspace*, within the *five-dimensional spacetime (bulk)*, referred to as *3-brane*. Many researches have been done around this new cosmology as e.g. [9]. We choose the *inflation in branes* [4], [6], [7], [8].

In this study, we consider that inflation might arise from the interaction potential between a 3-brane and anti-3-brane which are parallel and widely separated in five-dimensional Anti de Sitter space (*AdS₅*). The background is identical to that considered in the RS-I model [2].

The potential between the branes is given by [4]

$$V(\phi) \sim M_{5D}^4 \xi(\phi/M_{5D})(1 - e^{(-|\phi|/m)}). \quad (7)$$

The figures 1 and 2 show us the behaviours of the potential $V(\phi)$ and of the slow-roll parameters as a function of ϕ .

The spectral index n and its derivate $dn/d \ln k$ can be related to N , respectively, as

$$\frac{1}{2}(n - 1) = -\frac{1}{N}, \quad (8)$$

and

$$\frac{1}{2} \frac{dn}{d \ln k} = -\frac{1}{N^2}. \quad (9)$$

Setting $N = 70$ (as usually done in inflationary scenarios) leads to

$$n \approx 0,9714 \quad (10)$$

and

$$\frac{dn}{d \ln k} \approx -0,0004, \quad (11)$$

in excellent agreement with observational data from WMAP ($0.94 \leq n \leq 1.00$ and $-0.02 \leq dn/d \ln k \leq 0.02$, [5])

*Electronic address: michele@fma.if.usp.br




Figure 1: Behaviour of the potential $V(\phi)$ as a function ϕ .




Figure 2: Behaviour of the slow-roll parameters ϵ and η as a function of ϕ .

I. ACKNOWLEDGEMENT

This work was supported by CNPq.

-
- [1] Liddle, A. R. and Lyth, D. H., *Cosmological Inflation and Large-Scale Structure*, Cambridge University Press, Cambridge (2000).
 - [2] Randall, L. and Sundrum, R., *A Large Mass Hierarchy from a Small Extra Dimension*, Phys. Rev. Lett. 83: 3370 (1999).
 - [3] Randall, L. and Sundrum, R., *An Alternative to Compactification*, Phys. Rev. Lett. 83: 4690 (1999).
 - [4] Dvali, G. and Tye, S.-H Henry, *Brane Inflation*, Phys. Lett. B 450: 72-82 (1999).
 - [5] Peiris, H. V. et al., *First Year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Implications for Inflation*, Astrophys. J. Suppl. 148:213 (2003).
 - [6] Kachru, S., Kallosh, R. and Maldacena, J., *Towards Inflation in String Theory*, JCAP 0310:013 (2003).
 - [7] Kolda, C. and Lyth, D. H., *D-term Inflation and M-theory*, [hep-ph/98122343].
 - [8] Wang, B., Lin, C. Y. and Abdalla, E. *WMAP constrain on the P-term inflationary model*, Phys. Rev. D69:063507 (2004).
 - [9] Abdalla, E., Casali, A. G., Cuadros-Melgar, B. *Shortcuts in cosmological branes*, Int. J. Theor. Phys. 43:801-854 (2004).